

This is a reproduction of a library book that was digitized by Google as part of an ongoing effort to preserve the information in books and make it universally accessible.

GoogleTM books

<https://books.google.com>



Y3. A+7
22/USNRDL-TR-12
UNCLASSIFIED

F2

Copy No. 225


EFFECT OF REPETITIVE EXPOSURE TO GAMMA RAYS ON THE
HEMATOPOIETIC SYSTEM OF THE RAT

Research and Development Technical Report USNRDL-TR-12
NM 006 015

27 August 1954

by

S. J. Baum
D. J. Kimeldorf
E. M. Jacobsen



U.S. NAVAL RADIOLOGICAL DEFENSE LABORATORY

S A N F R A N C I S C O 2 4 C A L I F O R N I A

U. S. DEPOSITORY COPY

DOCUMENTS

UNIVERSITY OF MICHIGAN



3 9015 09516 3070

UNCLASSIFIED

Digitized by Google

Reproduction of this document in any form by other than activities of the Department of Defense is not authorized unless specifically approved by the Secretary of the Navy or the Chief of Naval Operations as appropriate.

U N C L A S S I F I E D

**EFFECT OF REPETITIVE EXPOSURE TO GAMMA RAYS ON THE
HEMATOPOIETIC SYSTEM OF THE RAT**

**Research and Development Technical Report USNRDL-TR-12
NM 006 015**

27 August 1954

by

**S. J. Baum
D. J. Kimeldorf
E. M. Jacobsen**

Health and Biology

**Technical Objective
AW-6**

**Physiology-Psychology Branch
D. J. Kimeldorf, Head**

**Biological and Medical Sciences Division
Captain A. R. Behnke, Jr. (MC) USN, Acting Head**

**Scientific Director
P. C. Tompkins**

**Director
Captain R. A. Hinnners, USN**

**U. S. NAVAL RADIOLOGICAL DEFENSE LABORATORY
San Francisco 24, California**

U N C L A S S I F I E D

Digitized by Google

ABSTRACT

Male rats were subjected to a series of one or more doses of 75 r gamma irradiation at seven day intervals between exposures. Hematological examinations were made following the last exposure in each series of irradiations. The maximum changes observed were, to a great extent, independent of the number of previous exposures, and appeared to reflect principally the effects of the last irradiation. These results were ascribed to the regenerative capacity of the hematopoietic system to recover during the seven day interval between radiation exposures. A discussion of these results and their correlation with the findings of other experiments is given.

SUMMARY

The Problem

To determine the effect of periodic exposure to low intensity gamma radiation upon hematopoiesis in the rat.

Findings

Rats were subjected to a series of one or more doses of 75 r gamma irradiation, at seven day intervals between exposures. Hematological examinations were made following the last exposure in each series of irradiations. The maximum changes observed were, to a great extent, independent of the number of previous exposures, and appeared to reflect principally the effects of the last irradiation. These results were ascribed to the regenerative capacity of the hematopoietic system to recover during the seven day interval between radiation exposures.

U N C L A S S I F I E D
=====

ADMINISTRATIVE INFORMATION

**This work was part of Project NM 006-015.04,
Phase 12, Technical Objective AW-6.**

**Acknowledgement: The authors wish to express
their appreciation to Mr. T. J. Castanera for his
assistance in the irradiation procedure.**

U N C L A S S I F I E D

REPORT OF INVESTIGATION

The hematological effects of acute ionizing radiations have been described extensively in recent review articles.^{1, 2, 4} The effects of prolonged exposure to low intensity radiations and the effects of dose fractionation or periodic radiation exposures are somewhat more difficult to interpret because of the complex interaction of destructive and regenerative factors involved.

When rats were exposed daily to low (7.7 r) doses of X rays relatively few changes were observed in the number of circulating blood cells, although the exposure was continued for 250 days.⁵ Mice subjected to 1.1, 4.4 or 8.8 r of gamma rays during eight hours daily⁶ for a one year period exhibited no significant erythrocyte changes. However, daily doses of 33.5 r of X rays⁵ have been demonstrated to be sufficient to cause a slight but distinct decrease in erythrocyte counts in rats. The dose-response relationship is somewhat species dependent. For example, a significant decrease in the erythrocyte estimate in guinea pigs exposed to 8.8 r gamma radiation for eight hours daily was observed by Lorenz, et al.⁷

Leukocytes are much more sensitive to ionizing radiation. Daily exposures to 7.7 r X irradiation caused a distinct decrease in the number of leukocytes.⁵

In the present experiment male rats were subjected to a series of one or more doses of 75 r gamma radiation at seven day intervals between exposures. Hematological examinations were made following the last exposure in each series of irradiations. This experimental design was selected in order to determine the effects of periodic exposure to sublethal doses of low intensity ionizing radiations upon recovery in the hematopoietic system as manifested by changes in the circulating blood cells. Presumably, observable effects would be the resultant of interaction between radiation damage caused by the periodic exposures and the forces of regeneration active during the interval.

EXPERIMENTAL DESIGN AND METHOD

Four hundred and thirty-two adult male Sprague-Dawley rats bred in the Laboratory colony were employed in this study. They were given food (Purina chow) and water ad lib. One half of the rats was distributed into six groups receiving different numbers of periodic exposures to gamma rays (75 r, 2 x 75 r, 3 x 75 r, 4 x 75 r, 6 x 75 r, and 8 x 75 r). These groups were divided further with respect to the six days of sacrifice (days 1, 5, 8, 14, 23, and 34), making a total of 36 irradiated groups. The remaining half of the animals comprised another set of 36 sham-irradiated groups which served as controls for the effects of radiation.

To facilitate experimental procedures and to control environmental variations, the total experimental pattern was repeated six times with one representative of each group for each replication. In each replication all animals were from litters born during the same week and were of similar body weight.

Animals were placed in small Lucite containers for irradiation and exposed to an isodose gamma radiation field (9.4 r/hr) for eight hours resulting in an air dose of 75 r for each irradiation. The radiation factors were: cobalt-60, approximately 1.3 Mev, 8.7 curies, point source. The controls were placed in the radiation exposure chambers and sham-irradiated.

On the days of sacrifice cardiac blood was drawn for the determination of the hematocrit, hemoglobin, and the erythrocyte, leukocyte and reticulocyte estimations. Standard Wright Stain blood smears were used for differential leukocyte counts. The mean corpuscular hemoglobin (M.C.H.), the mean corpuscular volume (M.C.V.), and the mean corpuscular hemoglobin concentration (M.C.H.C.), as listed in Table 1, are quotients for the ratios of hemoglobin/RBC count, hematocrit/RBC count, and hemoglobin/hematocrit, respectively.

Since each of the six irradiated animals in each group was matched with its own sham-irradiated control, the statistical analysis was based on differences between paired animals for each group. The probability tables for individual comparisons by the ranking methods of Wilcoxon⁸ were utilized for statistical evaluation of the data. Significant changes were referable to the probability level of $p \leq 0.05$.

RESULTS

A single exposure to 75 r low intensity gamma radiation during an eight hour period caused a significant decrease in the erythrocyte count of all groups by day 14 post radiation (Fig. 1). At that time a decrease of approximately 2×10^6 cells was observed. The erythrocyte count of rats receiving more than one dose of 75 r was always lower than that of the controls; however, the maximum depression was similar for all groups during the post-irradiation period regardless of the number of previous exposures.

Hemoglobin and hematocrit changes were in a similar direction to that of the erythrocyte count (Figs. 2, 3). The mean corpuscular hemoglobin value was larger for the irradiated rats on nearly all days of sacrifice regardless of the number of periodic exposures (Table 1). The maximum increase occurred on day 14 which was at the time of the lowest erythrocyte counts. The mean corpuscular volume of the irradiated rats was also larger than that of their respective controls (Table 1). The points of significant differences were nearly identical to those for the M.C.H. (Table 1). The mean corpuscular hemoglobin concentration was approximately 29 per cent for all groups and remained the same throughout the study (Table 1). There was no significant difference between irradiated rats and their controls.

The values for the reticulated erythrocytes of the exposed rats were low for the early days following the last exposure for each series of irradiations (Fig. 4). A recovery towards control counts was observed thereafter.

A decrease in the number of leukocytes was apparent on day one post irradiation (Fig. 5). The decrease was of similar magnitude for all irradiated groups regardless of the number of previous exposures. Recovery was initiated by day 5, and continued until day 34, at which point the values obtained from the irradiated rats were either normal or approaching it. The recovery in leukocyte counts was slightly more retarded in groups exposed more than once.

As can be observed from Fig. 5, the decrease in the total number of leukocytes was mainly due to a decline in lymphocytes. The neutrophil values obtained from irradiated rats were also always smaller than that of their controls (Fig. 6); however, their decline was less than that of the lymphocytes.

DISCUSSION

The anemia observed in animals irradiated with massive doses is a result of decreased hematopoiesis¹¹ and of a lowered capillary resistance which permits the escape of erythrocytes into the lymphatic system.¹⁴ However, the latter does not occur in animals irradiated with sublethal doses, as was demonstrated in recent experiments with rabbits.¹³ Presumably, changes in the number of erythrocytes in the rats of the present experiment which were subjected to low intensity radiation were caused by damage to the erythropoietic tissues. Such damage at low doses was observed by Bloom,¹⁰ who found that a dose of 80 r of X rays caused acute injury to the bone marrow as indicated by dead erythroblasts and increased nuclear debris. Indirect evidence was obtained by Hennessey and Huff.¹² They noticed that the iron uptake of rats subjected to 125 r of X rays was decreased about 40 per cent from the control values within the first five days post-irradiation. These authors stated that they were able to detect statistically significant depressions in tracer ion uptake in animals exposed to 5 and 25 r of X rays.

In the present experiment, the anemia of the irradiated rats was distinct, albeit slight and macrocytic. The cause of the macrocytic anemia is now known. Stearner, et al,⁵ observed this type of anemia in irradiated rats and Wintrobe⁹ reports such occurrences in human dial workers.

The outstanding result of this study was the extent of recovery apparently possible in the erythropoietic system during the seven day interval between periodic exposure to low intensity gamma irradiation, regardless of the number of previous exposures. Although differences attributable to previous periodic irradiations were observed one day after the last exposure, the extent of maximum red cell depression observed for all groups was similar and it occurred at about the same time (day 14). The latter fact suggests that the erythrocyte decline was, to a great extent, reflective of only the last exposure.

The remarkable ability for regeneration of the erythroid system has been observed in recent experiments. Bloom¹⁰ noticed in histological studies that erythroblasts, in spite of their great sensitivity to irradiation, were the first cells to regenerate. Valentine and Pearce¹⁵ irradiated (200 r, X rays) phlebotomized cats having a 40 per cent erythrocyte decrease and found the recovery from phlebotomy was only slightly inhibited. This procedure was repeated four times on six cats¹⁶ at four to six month intervals, and little evidence was obtained of any permanent damage to the erythropoietic system of these cats. The authors emphasized the fact that a dose of 200 r caused a profound insult to the hematological structure of cats, however, complete functional recovery occurred in the intervals between radiation exposures. It appears plausible that the relatively radioresistant stem cells of the marrow give first preference to the restoration of the impaired erythrocyte system.


U N C L A S S I F I E D

Erythropenia in the present study was preceded by a slight but significant reticulocytopenia. The latter returned to normal values after eight days. Ingram, et al.,³ reported such decreases in the reticulated cells of rats for single X-ray doses of 50 r and higher with recovery occurring approximately nine days after irradiation.

The early appearance of leukopenia in the present experiment was due to the short life span of leukocytes, primarily the lymphocytes. Inadequate production of new cells because of damaged myeloid and lymphoid precursors, therefore, will be observed within hours in the circulating blood. No additive destructive effect of previous exposures was detected, and in general recovery was good. However, regeneration was somewhat retarded in groups subjected to more than two doses of 75 r gamma radiation. This may indicate some residual influence upon recovery processes of the leukocyte system. It is of interest to note here that Valentine and Pearce¹⁵ observed a slower recovery of leukocytes in irradiated phlebotomized cats.

The results observed in the present study suggest that only slight, if any, cumulative effects are detectable in peripheral blood elements during the first 34 days after several (2-8) weekly exposures to low intensity gamma radiation. This is probably indicative of the recovery capacity of the hematopoietic tissues during the seven day interval between exposures.

Approved by:



A.R. BEHNKE, Jr.
Captain (MC) USN
Acting Head
Biological and Medical
Sciences Division

For the Scientific Director

U N C L A S S I F I E D

REFERENCES

1. Stelling, M. D., Effect of Radiation on the Blood. Radiography 18:221, 1952.
2. Furth, J. and Upton, A. C., Vertebrate Radiobiology: Histopathology and Carcinogenesis. Ann. Rev. Nuclear Sci. 3:303, 1953.
3. Ingram, M. and Mason, W. B., Effect of Acute Exposure to Roentgen Radiation on the Peripheral Blood of Experimental Animals. The University of Rochester Atomic Energy Project Unclassified Document UR-122, 1950.
4. Jacobson, L. O., Marks, E. K., Simmons, L. E., Hagen, C. W., Jr. and Zirkle, R. E., Effects of Total-Body X-Irradiation on Rabbits II. Hematological Effects, Chapter 8 in Biological Effects of External X and Gamma Radiation. McGraw-Hill Book Co., New York, 1954.
5. Stearner, S. P., Simmons, E. L. and Jacobson, L. O., The Effects of Total-Body X-Irradiation on the Peripheral Blood and Blood Forming Tissues of the Rat. Argonne National Laboratory Declassified Document ANL-4001 (MDDC-1319), 1947.
6. Spargo, B., Bloomfield, J. R., Glotzer, D. J., Grodon, E. and Nichols, O., Histological Effects of Long Continued Whole Body Gamma Irradiation of Mice. J. Nat. Cancer Inst. 12:615, 1951.
7. Lorenz, E., Heston, W. E., Jacobson, L. O., Eschenbrenner, A. B., Shimkin, M., Deringer, M. and Doninger, J., Biological Effects of Long Continued Whole Body Irradiation with Gamma Rays on Mice, Guinea Pigs, and Rabbits. (Part III). Metallurgical Laboratory Declassified Document MDDC-665, 1946.
8. Wilcoxon, F., Probability Tables for Individual Comparisons by Ranking Methods. Biometrics 3:3, 1947.
9. Wintrobe, M. M., Clinical Hematology. Lea and Febinger, Philadelphia, 1946.

10. Bloom, W., Histopathology of Irradiation. McGraw-Hill Book Co. Inc., New York, 1948.
11. Rosenthal, R. L., Pickering, B. J. and Goldschmidt, L. A Semi-Quantitative Study of Bone Marrow in Rats Following Total Body X-Irradiation. Blood 6:600, 1951.
12. Hennessy, T. G. and Huff, R. L., Depression of Tracer Iron Uptake Curve in Rat Erythrocytes Following Total Body X-Irradiation. Proc. Soc. Exptl. Biol. Med. 73:436, 1950.
13. Kahn, J. B. and Furth, J., The Pathogenesis of Post-Irradiation Anemia. Blood 7:404, 1952.
14. Ross, M. H., Furth, J. and Bigelow, R. R., Changes in Cellular Composition of the Lymph Caused by Ionizing Radiations. Blood 7:417, 1952.
15. Valentine, W. N. and Pearce, M. L., Studies on the Radiosensitivity of Bone Marrow I. The Relative Sensitivity of Erythroid and Myeloid Elements. Blood 7:1, 1952.
16. Valentine, W. N., Pearce, M. L. and Lawrence, J. S., Studies on the Radiosensitivity of Bone Marrow II. The Effect of Large, Repeated Whole Body Irradiation Exposure on Hematopoiesis. Blood 7:14, 1952.

TABLE 1
CALCULATED VALUES OF CELLULAR VOLUME,
HEMOGLOBIN AND HEMOGLOBIN CONCENTRATION

			Days of Sacrifice Post Irradiation					
	Dose (r)	Group	1	5	8	14	23	34
	75	C ¹	64.0	59.1	67.5	64.5	70.3	67.5
		I ²	65.6	<u>75.9</u>	70.7	<u>84.9</u>	76.4	78.9
Mean	2x75	C	63.5	66.3	69.7	56.9	61.7	68.7
		I	74.3	76.9	72.3	<u>85.6</u>	<u>79.1</u>	72.1
Corpuscular	3x75	C	68.9	73.2	66.6	63.4	63.4	68.7
Volume		I	74.7	<u>82.7</u>	75.5	<u>83.7</u>	<u>83.4</u>	82.7
cubic	4x75	C	63.1	68.4	63.4	61.2	60.9	64.1
		I	73.1	76.2	<u>82.9</u>	<u>85.2</u>	<u>76.3</u>	68.4
microns	6x75	C	71.3	69.9	70.7	71.6	67.9	71.4
		I	71.5	<u>82.6</u>	74.9	<u>85.3</u>	74.5	68.7
	8x75	C	66.8	75.4	65.8	64.4	67.3	67.7
		I	<u>75.2</u>	80.8	<u>83.1</u>	<u>88.9</u>	<u>78.2</u>	<u>78.9</u>
	75	C	19.2	17.6	20.1	18.8	20.2	19.4
		I	19.2	<u>23.3</u>	20.9	<u>26.2</u>	22.7	24.4
Mean	2x75	C	19.0	19.5	20.1	18.4	18.1	20.0
		I	22.0	23.2	21.6	<u>25.3</u>	22.4	20.1
Corpuscular	3x75	C	19.8	21.4	19.4	18.5	19.2	19.8
Hemoglobin		I	22.4	<u>24.0</u>	22.3	<u>25.3</u>	<u>23.5</u>	23.5
micro	4x75	C	18.1	19.7	18.2	19.6	18.0	18.0
		I	20.1	22.0	<u>23.9</u>	<u>24.5</u>	<u>21.7</u>	19.5
micrograms	6x75	C	19.9	19.6	20.2	21.4	19.7	20.6
		I	21.1	<u>25.6</u>	21.8	<u>25.0</u>	<u>22.8</u>	19.6
	8x75	C	18.9	21.4	18.4	19.7	19.6	19.4
		I	<u>22.1</u>	24.2	<u>25.7</u>	<u>26.7</u>	<u>22.9</u>	23.4
	75	C	29.9	29.9	29.9	29.2	28.8	28.7
		I	29.5	30.6	29.7	31.5	29.7	30.8
Mean	2x75	C	30.0	29.4	29.6	30.3	29.4	29.1
		I	29.7	30.0	30.0	29.5	28.5	28.9
Corpuscular	3x75	C	28.8	29.2	29.2	29.2	30.4	28.8
Hemoglobin		I	30.8	29.1	29.5	30.3	28.1	28.6
Concentration	4x75	C	28.8	28.8	28.6	30.6	30.1	28.1
		I	28.3	28.9	28.6	28.8	28.4	30.0
per cent	6x75	C	28.0	28.0	28.2	29.9	28.8	28.8
		I	29.9	32.0	29.1	30.5	30.7	28.6
	8x75	C	28.3	28.3	28.0	28.9	29.2	28.7
		I	29.3	30.1	31.0	30.1	29.5	29.1

1. C = Controls

2. I = Irradiated Rats

3. Underlined mean values of pairs differ significantly, $P \leq 0.05$.




Fig. 1 Erythrocyte Values of Rats Subjected to Various Numbers of 75 r Doses of Gamma Radiation. Significant (5 per cent level) differences between mean values are indicated by asterisks.




Fig. 2 Hemoglobin Values of Rats Subjected to Various Numbers of 75 r Doses of Gamma Radiation. Significant (5 per cent level) differences between mean values are indicated by asterisks.




Fig. 3 Hematocrit Values of Rats Subjected to Various Numbers of 75 r Doses of Gamma Radiation. Significant (5 per cent level) differences between mean values are indicated by asterisks.




Fig. 4 Reticulocyte Values of Rats Subjected to Various Numbers of 75 r Doses of Gamma Radiation. Significant (5 per cent level) differences between mean values are indicated by asterisks.




Fig. 5 Leukocyte and Lymphocyte Values of Rats Subjected to Various Numbers of 75 r Doses of Gamma Radiation. Significant (5 per cent level) differences between mean values are indicated by asterisks.




Fig. 6 Neutrophil Values of Rats Subjected to Various Numbers of 75 r Doses of Gamma Radiation. Significant (5 per cent level) differences between mean values are indicated by asterisks.

U N C L A S S I F I E D

DISTRIBUTION

COPIES

NAVY

1-9 Chief, Bureau of Ships (Code 324)
10-11 Chief, Bureau of Ships (Code 366, for forwarding)
12 Chief, Bureau of Medicine and Surgery
13 Chief, Bureau of Medicine and Surgery (for forwarding)
14 Chief, Bureau of Aeronautics (Code AE54)
15 Chief, Bureau of Supplies and Accounts (Code W)
16-17 Chief, Bureau of Yards and Docks (D-440)
18 Chief of Naval Research
19 Chief of Naval Operations (Op-36)
20 Commander, New York Naval Shipyard (Material Lab.)
21-23 Director, Naval Research Laboratory (Code 2021)
24 Naval Medical Research Institute
25 CO, Naval Unit, Army Chemical Center
26 CO, Naval Unit, CmlC Training Command
27 CO, U.S. Naval Civil Engineering (Res. and Eval. Lab.)
28 U.S. Naval School (CEC Officers)
29 CO, Naval Air Material Center, Philadelphia
30 CO, Naval Air Development Center
31 CO, Naval Schools Command, Treasure Island
32 CO, Naval Damage Control Training Center, Philadelphia
33 CO, Fleet Training Center, Norfolk
34 CO, Fleet Training Center, San Diego
35 Commander Training Command, Pacific Fleet
36 Office of Patent Counsel, Mare Island
37 Commander Air Force, Atlantic Fleet (Code 16F)
38 Commandant, U.S. Marine Corps
39-40 Commandant, Marine Corps Schools, Quantico

ARMY

41 Chief of Engineers, Department of the Army (ENGEB, Dhein)
42-44 CO, CmlC Chemical and Radiological Laboratories (Library)
45-46 CG, CmlC Res. and Eng. Command, ACmlC, Maryland

U N C L A S S I F I E D

47 CG, Chemical Corps Materiel Command
48 Radiological Division, CmlC Chemical and Radiological Lab.
49 CO, Chemical Corps Medical Laboratories
50 Chemical Corps Training Command (Library)
51 Office of Chief Signal Officer (SIGGG-P)
52 CO, Office of Ordnance Research (ORDOR-1A)
53 CO, Army Medical Research Laboratory, Fort Knox
54-55 Army Medical Research Laboratory, Fort Knox
56 Army Medical Service Graduate School
57 CG, Quartermaster Res. and Dev. Center
58 CG, Quartermaster Res. and Dev. Command
59 Operations Research Office
60 CO, Engineer Res. and Dev. Lab.
61 Dugway Proving Ground
62 The Surgeon General
63 Director, Evans Signal Laboratory (Nucleonics Section)
64 CO, Transportation Res. and Dev. Command, Fort Eustis
65 Tokyo Army Hospital
66 Watertown Arsenal
67 Frankford Arsenal

AIR FORCE

68 Directorate of Research and Development (AFDRD-HF)
69 CO, Air Materiel Command (MCMTM)
70 CG, Wright Air Development Center (WCOES)
71 CG, Wright Air Development Center (WCRDO)
72-74 CG, Wright Air Development Center (WCRTY)
75 CG, Wright Air Development Center (WCRTH-1)
76 CG, Air Res. and Dev. Command (RDTDA)
77 CG, Air Res. and Dev. Command (RDTRH)
78 USAF, School of Aviation Medicine
79 USAF, School of Aviation Medicine (Brooks)
80 CG, Strategic Air Command, Offutt Air Force Base (IGABD)
81 CG, Strategic Air Command (Operations Analysis Office)
82 CG, Kirtland Air Force Base (Library)
83 Director, Air University Library, Maxwell Air Force Base
(CR-4030)
84-85 CG, Department of Armament Training, Lowry Air Force
Base
86 CG, Cambridge Research Center (CRHK)

OTHER DOD ACTIVITIES

87 Chief, Armed Forces Special Weapons Project
88 AFSWP, Technical Training Group, Sandia Base
89 CG, AFSWP, Hq., Field Command, Sandia Base

U N C L A S S I F I E D

90 Assistant Secretary of Defense (Res. and Dev.)
91 Armed Forces Medical Library
92-96 Armed Services Technical Information Agency
97 Armed Forces Institute of Pathology

AEC ACTIVITIES AND OTHERS

98-100 Argonne Cancer Research Hospital
101-106 Argonne National Laboratory
107-108 Atomic Bomb Casualty Commission (APO-182)
109 Atomic Bomb Casualty Commission, Washington
110-112 AEC, Washington
113 Atomic Energy of Canada, Ltd.
114 Australian Atomic Energy Commission
115 Battelle Memorial Institute
116-118 Belgium, Union Miniere du Haut Katanga
119-122 Brookhaven National Laboratory
123 Brush Beryllium Company
124-126 Canadian Joint Staff
127 Carbide and Carbon Chemicals Company (C-31 Plant)
128-129 Carbide and Carbon Chemicals Company (K-25 Plant)
130-135 Carbide and Carbon Chemicals Company (ORNL)
136 Centre d'Etudes pour les Applications de l'Energie Nucleaire
137 Centro Italian Studi Esperienze
138-141 Chalk River Project, Canada
142 Chicago Patent Group
143 Columbia University (Failla)
144 Columbia University (Hassialis)
145 Commonwealth X-Ray and Radium Laboratory
146 Consolidated Vultee Aircraft Corporation
147 Division of Raw Materials, Denver
148-149 Dow Chemical Company, Midland
150 Dow Chemical Company, Rocky Flats
151-152 duPont Company, Savannah
153 duPont Company, Wilmington
154 General Electric Company (ANPP)
155-156 General Electric Company, Richland
157 Goodyear Atomic Corporation
158 Harshaw Chemical Corporation
159 Iowa State College
160 Kaiser Engineers
161-162 Knolls Atomic Power Laboratory
163 Lockheed Aircraft Corporation
164-165 Los Alamos Scientific Laboratory
166 Mallinckrodt Chemical Works
167 Massachusetts Institute of Technology (Hardy)
168 Massachusetts Institute of Technology (Kaufmann)

U N C L A S S I F I E D

169 Mound Library
170 National Bureau of Standards (Taylor)
171 National Bureau of Standards (Library)
172 National Lead Company of Ohio
173-175 National Research Council, Ottawa
176 New Brunswick Laboratory
177 Newport News Shipbuilding and Drydock Company
178 New York Operations Office
179 North American Aviation, Inc.
180 Nuclear Development Associates, Inc.
181 Oak Ridge Institute of Nuclear Studies
182 Patent Branch, Washington
183-188 Phillips Petroleum Company
189-191 Public Health Service, Washington
192 RAND Corporation
193 Tokyo University
194 United Aircraft Corporation
195-209 United Kingdom Scientific Mission
210 U.S. Geological Survey (Butler)
211 UCLA Medical Research Laboratory
212 University of California Medical Center
213-215 University of California Radiation Laboratory, Berkeley
216-217 University of California Radiation Laboratory, Livermore
218 University of Chicago Radiation Laboratory
219-220 University of Michigan (Gomberg)
221-222 University of Rochester (Technical Report Unit)
223 University of Tennessee (Comar)
224 University of Utah (Bowers)
225 University of Washington, Applied Fisheries Lab.
226 Vitro Corporation of America
227-230 Western Reserve University
231-232 Westinghouse Electric Company
233-257 Technical Information Service, Oak Ridge

USNRDL

258-275 USNRDL, Technical Information Division

DATE ISSUED: 2 November 1954

<p>Naval Radiological Defense Laboratory. USNRDL-TR-12.</p> <p>EFFECT OF REPETITIVE EXPOSURE TO GAMMA RAYS ON THE HEMATOPOIETIC SYSTEM OF THE RAT, by S.J. Baum, D. J. Kimeldorf, and E. M. Jacobsen. 27 Aug. 1954. iv, 18 p. diagrs. (16 refs.) UNCLASSIFIED</p> <p>Male rats were exposed one or more times to 75 r gamma at seven day intervals. Hematological examinations after the last exposure indicated that the maximum changes were largely independent of the number of exposures and reflected mostly the effects of the last irradiation. These results are ascribed to the capacity of the hematopoietic system to recover during the seven day intervals.</p>	<p>1. Gamma radiation - Physiological effects 2. Blood forming organs - Radiation injuries I. Baum, S.J. II. Kimeldorf, D.J. III. Jacobsen, E. M. IV. Title V. NM 006 015</p>	<p>UNCLASSIFIED</p>	<p>Naval Radiological Defense Laboratory. USNRDL-TR-12.</p> <p>EFFECT OF REPETITIVE EXPOSURE TO GAMMA RAYS ON THE HEMATOPOIETIC SYSTEM OF THE RAT, by S.J. Baum, D.J. Kimeldorf, and E. M. Jacobsen. 27 Aug. 1954. iv, 18 p. diagrs. (16 refs.) UNCLASSIFIED</p> <p>Male rats were exposed one or more times to 75 r gamma at seven day intervals. Hematological examinations after the last exposure indicated that the maximum changes were largely independent of the number of exposures and reflected mostly the effects of the last irradiation. These results are ascribed to the capacity of the hematopoietic system to recover during the seven day intervals.</p>	<p>1. Gamma radiation - Physiological effects 2. Blood forming organs - Radiation injuries I. Baum, S.J. II. Kimeldorf, D.J. III. Jacobsen, E. M. IV. Title V. NM 006 015</p>	<p>UNCLASSIFIED</p>
<p>Naval Radiological Defense Laboratory. USNRDL-TR-12.</p> <p>EFFECT OF REPETITIVE EXPOSURE TO GAMMA RAYS ON THE HEMATOPOIETIC SYSTEM OF THE RAT, by S.J. Baum, D. J. Kimeldorf, and E. M. Jacobsen. 27 Aug. 1954. iv, 18 p. diagrs. (16 refs.) UNCLASSIFIED</p> <p>Male rats were exposed one or more times to 75 r gamma at seven day intervals. Hematological examinations after the last exposure indicated that the maximum changes were largely independent of the number of exposures and reflected mostly the effects of the last irradiation. These results are ascribed to the capacity of the hematopoietic system to recover during the seven day intervals.</p>	<p>1. Gamma radiation - Physiological effects 2. Blood forming organs - Radiation injuries I. Baum, S.J. II. Kimeldorf, D.J. III. Jacobsen, E. M. IV. Title V. NM 006 015</p>	<p>UNCLASSIFIED</p>	<p>Naval Radiological Defense Laboratory. USNRDL-TR-12.</p> <p>EFFECT OF REPETITIVE EXPOSURE TO GAMMA RAYS ON THE HEMATOPOIETIC SYSTEM OF THE RAT, by S.J. Baum, D.J. Kimeldorf, and E. M. Jacobsen. 27 Aug. 1954. iv, 18 p. diagrs. (16 refs.) UNCLASSIFIED</p> <p>Male rats were exposed one or more times to 75 r gamma at seven day intervals. Hematological examinations after the last exposure indicated that the maximum changes were largely independent of the number of exposures and reflected mostly the effects of the last irradiation. These results are ascribed to the capacity of the hematopoietic system to recover during the seven day intervals.</p>	<p>1. Gamma radiation - Physiological effects 2. Blood forming organs - Radiation injuries I. Baum, S.J. II. Kimeldorf, D.J. III. Jacobsen, E. M. IV. Title V. NM 006 015</p>	<p>UNCLASSIFIED</p>

<p>Naval Radiological Defense Laboratory. USNRDL-TR-12.</p> <p>EFFECT OF REPETITIVE EXPOSURE TO GAMMA RAYS ON THE HEMATOPOIETIC SYSTEM OF THE RAT, by S.J. Baum, D. J. Kimeldorf, and E. M. Jacobsen. 27 Aug. 1954. iv, 18 p. diags. (16 refs.) UNCLASSIFIED</p> <p>Male rats were exposed one or more times to 75 r gamma at seven day intervals. Hematological examinations after the last exposure indicated that the maximum changes were largely independent of the number of exposures and reflected mostly the effects of the last irradiation. These results are ascribed to the capacity of the hematopoietic system to recover during the seven day intervals.</p>	<p>1. Gamma radiation - Physiological effects</p> <p>2. Blood forming organs - Radiation injuries</p> <p>I. Baum, S.J.</p> <p>II. Kimeldorf, D.J.</p> <p>III. Jacobsen, E.M.</p> <p>IV. Title</p> <p>V. NM 006 015</p> <p>UNCLASSIFIED</p>
<p>Naval Radiological Defense Laboratory. USNRDL-TR-12.</p> <p>EFFECT OF REPETITIVE EXPOSURE TO GAMMA RAYS ON THE HEMATOPOIETIC SYSTEM OF THE RAT, by S.J. Baum, D.J. Kimeldorf, and E. M. Jacobsen. 27 Aug. 1954. iv, 18 p. diags. (16 refs.) UNCLASSIFIED</p> <p>Male rats were exposed one or more times to 75 r gamma at seven day intervals. Hematological examinations after the last exposure indicated that the maximum changes were largely independent of the number of exposures and reflected mostly the effect of the last irradiation. These results are ascribed to the capacity of the hematopoietic system to recover during the seven day intervals.</p>	<p>1. Gamma radiation - Physiological effects</p> <p>2. Blood forming organs - Radiation injuries</p> <p>I. Baum, S.J.</p> <p>II. Kimeldorf, D.J.</p> <p>III. Jacobsen, E.M.</p> <p>IV. Title</p> <p>V. NM 006 015</p> <p>UNCLASSIFIED</p>
<p>Naval Radiological Defense Laboratory. USNRDL-TR-12.</p> <p>EFFECT OF REPETITIVE EXPOSURE TO GAMMA RAYS ON THE HEMATOPOIETIC SYSTEM OF THE RAT, by S.J. Baum, D. J. Kimeldorf, and E. M. Jacobsen. 27 Aug. 1954. iv, 18 p. diags. (16 refs.) UNCLASSIFIED</p> <p>Male rats were exposed one or more times to 75 r gamma at seven day intervals. Hematological examinations after the last exposure indicated that the maximum changes were largely independent of the number of exposures and reflected mostly the effects of the last irradiation. These results are ascribed to the capacity of the hematopoietic system to recover during the seven day intervals.</p>	<p>1. Gamma radiation - Physiological effects</p> <p>2. Blood forming organs - Radiation injuries</p> <p>I. Baum, S.J.</p> <p>II. Kimeldorf, D.J.</p> <p>III. Jacobsen, E.M.</p> <p>IV. Title</p> <p>V. NM 006 015</p> <p>UNCLASSIFIED</p>
<p>Naval Radiological Defense Laboratory. USNRDL-TR-12.</p> <p>EFFECT OF REPETITIVE EXPOSURE TO GAMMA RAYS ON THE HEMATOPOIETIC SYSTEM OF THE RAT, by S.J. Baum, D.J. Kimeldorf, and E. M. Jacobsen. 27 Aug. 1954. iv, 18 p. diags. (16 refs.) UNCLASSIFIED</p> <p>Male rats were exposed one or more times to 75 r gamma at seven day intervals. Hematological examinations after the last exposure indicated that the maximum changes were largely independent of the number of exposures and reflected mostly the effects of the last irradiation. These results are ascribed to the capacity of the hematopoietic system to recover during the seven day intervals.</p>	<p>1. Gamma radiation - Physiological effects</p> <p>2. Blood forming organs - Radiation injuries</p> <p>I. Baum, S.J.</p> <p>II. Kimeldorf, D.J.</p> <p>III. Jacobsen, E.M.</p> <p>IV. Title</p> <p>V. NM 006 015</p> <p>UNCLASSIFIED</p>

UNCLASSIFIED